



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Comparing the two results we see that the cone empties in  $\frac{8}{15}$  of the time it takes the cylinder, or the cylinder takes  $1\frac{1}{2}$  as long as the cone to empty. The minus sign is prefixed because  $x$  decreases as  $t$  increases.

Also solved by G. B. M. ZERR, ELMER SCHUYLER, J. SCHEFFER, and J. C. NAGLE.

NOTE. In reference to problem 63, Dr. Arnold Emch says: "I had the problem solved by my class in graphic statics by a purely graphical method and the following values (approximations) were obtained:  $\angle ABE=46^\circ$ ,  $\angle BAD=56^\circ$ , tension in  $BE=56.8$ , tension in  $AD=71$ . This shows that the solution in the MONTHLY is correct."

### DIOPHANTINE ANALYSIS.

71. Proposed by A. H. BELL, Hillsboro, Ill.

Find five numbers such that the product of any two plus 1 will equal a square.

III. Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

By using  $(s-1)^2$ , the denominator of Euler's fifth number, where  $s=4n(n-1)(n+1)[4n(2n-1)(2n+1)]$ , I have found five numbers in terms of  $n$ :  $x=n-1$ ,  $y=n+1$ ,  $z=4n$ ,  $w=4n(2n-1)(2n+1)$ , and

$$v = \frac{4n(2n-1)(2n+1)[2n(2n-1)-1][2n(2n+1)-1](8n^2-1)}{\{4n(n-1)(n+1)[4n(2n-1)(2n+1)]-1\}^2}.$$

The numerator of  $v$  is four times the product of the roots of the six squares  $xy+1$ ,  $xz+1$ ,  $yz+1$ ,  $xw+1$ ,  $yw+1$ , and  $zw+1$ .

The denominator of  $v$  is the square of  $(xyzw-1)$ .

Take  $n=1, 2, 3, 4, 5, 6$ , etc. We then obtain the following sets of five numbers:

$$\begin{array}{lllll} 0, & 2, & 4, & 12, & 420; \\ 1, & 3, & 8, & 120, & \frac{777480}{(2879)^2}; \\ 2, & 4, & 12, & 420, & \frac{35455980}{(40319)^2}; \\ 3, & 5, & 16, & 1008, & \frac{499902480}{(241919)^2}; \\ 4, & 6, & 20, & 1980, & \frac{3822388020}{(950399)^2}; \\ 5, & 7, & 24, & 3432, & \frac{20000100120}{(2882879)^2}; \text{ etc., etc., etc.} \end{array}$$

We also find

$$xv+1=\frac{\{(2n-1)(2n+1)[2n(2n-1)-1][2n(2n+1)-1]-2n(8n^2-1)\}^2}{(s-1)^2};$$

$$yv+1=\frac{\{(2n-1)(2n+1)[2n(2n-1)-1][2n(2n+1)-1]+2n(8n^2-1)\}^2}{(s-1)^2};$$

$$zv+1=\frac{\{16n^4(2n-1)(2n+1)-(8n^2-1)\}^2}{(s-1)^2}; \text{ and}$$

$$wv+1=\frac{\{24n^2(2n^2-1)(2n-1)(2n+1)-1\}^2}{(s-1)^2}.$$

74. Proposed by O. W. ANTHONY, M. Sc., Instructor in Mathematics, Boys' High School, New York City.

Solve  $x^2+y^2=\square$ ,  $z^2+w^2=\square$ ,  $y^2+w^2=\square$ .

I. Solution by M. A. GRUBER, A. M., War Department, Washington. D. C.

Take any two integral equations in which the sum of two squares equals a square, as

$$a^2+b^2=c^2, \text{ and } a_1^2+b_1^2=c_1^2.$$

Multiply the terms of the first equation by the first term and the second term, respectively, of the second equation. Also multiply the terms of the second equation by the first term and the second term, respectively, of the first equation. We then have

$$(aa_1)^2+(a_1b)^2=(a_1c)^2 \dots (1),$$

$$(ab_1)^2+(bb_1)^2=(b_1c)^2 \dots (2),$$

$$(aa_1)^2+(ab_1)^2=(ac_1)^2 \dots (3),$$

$$(a_1b)^2+(bb_1)^2=(bc_1)^2 \dots (4).$$

Now put  $x=aa_1$ ,  $y=a_1b$ ,  $z=ab_1$ , and  $w=bb_1$ ; then equations (1), (2), and (4) are the three required by the problem, there being added, in the solution,  $x^2+z^2=\square \dots (3)$ .

By means of the formula  $(2mn)^2+(m^2-n^2)^2=(m^2+n^2)^2$ , find a few integral numerical equations.

Take  $m=2$ ,  $n=1$ ; then  $4^2+3^2=5^2 \dots (1)$ .

Take  $m=3$ ,  $n=2$ ; then  $12^2+5^2=13^2 \dots (2)$ .

Take  $m=4$ ,  $n=1$ ; then  $8^2+15^2=17^2 \dots (3)$ .

Take  $m=5$ ,  $n=2$ ; then  $20^2+21^2=29^2 \dots (4)$ , etc.

From (1) and (2),  $x=48$ ,  $y=36$ ,  $z=20$ ,  $w=15$ .

From (1) and (3),  $x=32$ ,  $y=24$ ,  $z=60$ ,  $w=45$ .

From (1) and (4),  $x=80$ ,  $y=60$ ,  $z=84$ ,  $w=63$ .

From (2) and (3),  $x=96$ ,  $y=40$ ,  $z=180$ ,  $w=75$ , etc.